

Considerations for Pediatric Craniofacial Surgeons During the COVID-19 Outbreak

Anna Schoenbrunner, MD,* Benjamin Sarac, BS,[†]
Amanda Gosman, MD,[‡] and Jeffrey E. Janis, MD*

The Coronavirus Disease 2019 (COVID-19) outbreak was declared a pandemic by the World Health Organization (WHO) on March 11, 2020. One week later, the Centers for Medicare and Medicaid Services (CMS) issued a statement that “all elective, nonessential medical, surgical, and dental procedures be delayed” during the COVID-19 outbreak to minimize spread of the SARS-CoV-2 virus and preserve personal protective equipment (PPE).¹ The CMS recommendations created a tiered framework to guide surgeons and healthcare systems in their determination of which procedures are elective based on acuity, risk of progression, and symptomatology. The guidelines also take into account staffing support, ventilator availability, and patient comorbidities. The CMS report spurred action at the state level, with similar recommendations issued regarding postponement of elective surgeries and procedures. This report presents recommendations issued by state governing bodies as of 5PM EDT on March 24, 2020. Thirty states and the District of Columbia provide recommendations or orders regarding the postponement of elective procedures. As of this time, the only state to mention consideration of patient age is Arizona, which recommends surgeons “consider the health and age of the patient.”²

The pediatric population has been largely spared during the COVID-19 outbreak, with incidence of COVID-19 infection of less than 1% in children under 10 years of age and less than 2% in children under 19 years of age.^{3–6} However, children have been found to be asymptomatic carriers of SARS-CoV-2 with reports of transmissions from asymptomatic children to adults.^{7,8} Of special note, aerosolizing procedures increase transmission risk of COVID-19; craniofacial surgeons face high risk of COVID-19 infection due to exposure risks of operating in or near the oral cavity.^{9,10} Current recommendations for PPE use during surgical procedures on suspected or confirmed COVID-19 patients suggest the use of either a N95 or powered air-purifying respirator (PAPR).¹¹ Clinical stability

From the *Department of Plastic and Reconstructive Surgery, The Ohio State University Wexner Medical Center; [†]The Ohio State University College of Medicine, Columbus, OH; and [‡]Division of Plastic Surgery, University of California, San Diego, San Diego, CA.

Received March 27, 2020.

Accepted for publication March 28, 2020.

Address correspondence and reprint requests to Jeffrey E. Janis, MD, 915 Olentangy River Road, Suite 2140, Columbus, OH 43212; E-mail: Jeffrey.janis@osumc.edu

Information presented is intended to educate the reader on the variety of guidance published by state governments regarding surgical procedures and should not be used as a substitute for each reader's respective local or state guidance. Given the rapidly evolving nature of the COVID-19 pandemic, official guidance may have changed during data collection or by the time you are reading this.

The authors report no conflicts of interest.

Supplemental digital contents are available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.jcraniofacialsurgery.com).

Copyright © 2020 by Mutaz B. Habal, MD

ISSN: 1049-2275

DOI: 10.1097/SCS.00000000000006565

of a patient with active COVID-19 infection and risk of transmission to healthcare staff must be carefully considered before surgical intervention.

The pediatric surgical population is unique for several reasons; most notably the time-sensitive nature of many pediatric interventions. This poses a unique challenge when applying guidelines aimed at an adult population; no state recommendations address the nuances of pediatric surgical practice. We discuss common craniofacial procedures and relate these to the CMS tiered framework. We make special note of cases in which the time-sensitive nature of a condition may prompt more urgent surgical intervention. We reference state recommendations as applicable. Recommendations contained in this report (Supplemental Digital Content, Table 1, <http://links.lww.com/SCS/B428>) should serve only as general guidance; craniofacial surgeons must individualize treatment recommendations based on their patients' unique circumstances as well as their institutional and state policies.

CLEFT LIP/PALATE AND ASSOCIATED ANCILLARY PROCEDURES

Primary cleft lip and palate repair is a low acuity surgery performed in a healthy patient, making this a Tier 1a procedure; CMS recommends these surgeries be postponed. However, cleft palate repair performed after the age of 12 months is associated with worse speech outcomes; for this reason, cleft palate repair is usually favored to be performed prior to 12 months of age.^{12,13} Cleft lip and palate revisions, including lip revision, tip rhinoplasty, and speech surgery, are likewise Tier 1a procedures and should be postponed. Alveolar bone grafting is a time-sensitive procedure as this must be timed with eruption of the permanent canines; this timing should be taken into account when scheduling the intervention.

No state guidelines specifically mention cleft lip or palate repair or associated procedures.

ORTHOGNATHIC SURGERY

Orthognathic surgery is a low acuity surgery generally performed in a healthy patient, making this a Tier 1a procedure. Though timing of orthognathic surgery is dependent on orthodontic treatment and skeletal maturity, postponement of orthognathic surgery does not interfere with orthodontic treatment.

No state guidelines specifically mention orthognathic surgery; however, 12 states and the District of Columbia specifically mention that elective dental procedures be postponed.^{2,14–24}

MANDIBULAR DISTRACTION OSTEOGENESIS/TONGUE LIP ADHESION

Mandibular distraction osteogenesis (MDO) or tongue lip adhesion (TLA) is typically performed for patients with airway obstruction due to retrognathia and glossoptosis in Pierre Robin Sequence. In situations where MDO or TLA is performed to avoid intubation or tracheostomy, the procedure would be classified as Tier 3a and should not be postponed. In situations where the patient is healthy in an outpatient clinic setting, MDO or TLA is classified as Tier 1a; however, the age of the patient must be taken into account to determine importance of timing of the procedure.

No state guidelines specifically mention MDO or TLA procedures.

CRANIOSYNOSTOSIS

Craniosynostosis poses unique challenges in management, as intervention is time sensitive. All management decisions should be made in conjunction with neurosurgery colleagues. For patients who present early and who are candidates for minimally invasive procedures, intervention is favored prior to 4 months of age; these patients are classified as Tier 2a (consider postponing) or Tier 3a

(do not postpone). However, postponing an intervention beyond 4 months of age may make these patients ineligible for minimally invasive approaches, thereby putting patients at higher risk of complication.^{25,26} In such instances, surgical intervention may be considered prior to 4 months of age, based on institutional practices.

For patients who are outside of the window for minimally invasive procedures, intervention for open cranial vault remodeling remains time sensitive as patients who undergo the procedure over 12 months of age can have decreased rates of reossification and higher complication rates.^{27,28} Based on the CMS classification, these patients are classified as Tier 2a; however, due to the risk of complications and decreased reossification rates, open cranial vault remodeling is ideally performed prior to 12 months of age. In situations where patients present with findings of elevated intracranial pressure, acute intervention is required and these patients are classified as Tier 3a; this intervention should not be postponed.

No state guidelines specifically mention craniosynostosis; however, 5 states mention that elective cases should only be postponed if they can safely be done so within a period of 30 days to 3 months.^{21,23,29–31} As craniosynostosis cases are time sensitive, such considerations may apply.

FACIAL TRAUMA

Facial trauma encompasses a range of clinical scenarios, each with unique considerations. Many facial fractures are nonoperative and require no more than symptomatic management and precautions. For operative fractures, the risk for airway compromise, health of the patient, and acuity of intervention must be assessed; interventions for facial fractures range from Tier 1a for interventions based on cosmetic concerns and Tier 3a for highly symptomatic patients. Facial nerve repair for acute facial nerve injury after trauma is classified as Tier 3b and should not be postponed.

No state guidelines specifically mention facial trauma. However, Minnesota's recommendations specifically state that if there is "threat of permanent dysfunction of an extremity or organ system, including teeth and jaws" that such an intervention is considered nonelective.²⁰

OTHER CRANIOFACIAL PROCEDURES

Craniofacial surgeons perform a variety of other procedures in their practice not specifically addressed above. Surgeons are advised to weigh the acuity of each procedure and health of the patient against the risk of SARS-CoV-2 exposure and impact on the health system. Most procedures will be low acuity in healthy patients (Tier 1a) and should be postponed.

Procedures involving excision of benign or malignant tumors should be carefully considered; national oncology organizations have recommended postponing oncologic resection procedures for cancers that are at low risk for progression or metastases; the length of postponement varies based on the individual guideline and health system resources.^{32,33} Most oncologic resection procedures performed by craniofacial surgeons are for tumors with low risk of progression or metastasis; these are classified as 2a and postponement should be considered. However, those tumors with high risk of progression or metastasis are classified as Tier 3a (do not postpone).

Nine states specifically mention cancer in their guidelines; these guidelines make exceptions for cancer cases and cases in which there is a risk of metastasis or progression of staging of disease.^{2,14,17,20,21,23,31,34,35}

CONCLUSION

Craniofacial surgeons must balance the risks of postponing a surgical procedure with the risk of exposure to the child and

healthcare staff and risk of developmental delay against delaying the procedure during the COVID-19 outbreak. The CMS guidelines and state-specific guidelines provide guidance to craniofacial surgeons; however, the time-sensitive nature of many pediatric craniofacial interventions must be considered.

REFERENCES

- Centers for Medicare and Medicaid Services. CMS Adult Elective Surgery and Procedures Recommendations: Limit all non-essential planned surgeries and procedures, including dental, until further notice [press release]. Available at: <https://www.cms.gov/files/document/covid-elective-surgery-recommendations.pdf>. Accessed 2020
- State of Arizona. Delaying Elective Surgeries to Conserve Personal Protective Equipment to Test and Treat Patients with COVID-19. Available at: <https://azgovernor.gov/file/34242/download?token=13JvjX7I>. Accessed March 21, 2020
- Lee PI, Hu YL, Chen PY, et al. Are children less susceptible to COVID-19? *J Microbiol Immunol Infect* [published ahead of print February 25, 2020] doi: 10.1016/j.jmii.2020.02.011
- Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* [published ahead of print February 7, 2020] doi: 10.1001/jama.2020.1585
- Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497–506
- Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med* 2020;382:1199–1207
- Bai Y, Yao L, Wei T, et al. Presumed asymptomatic carrier transmission of COVID-19. *JAMA* [published ahead of print February 21, 2020] doi: 10.1001/jama.2020.2565
- Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 2020;395:514–523
- Meng L, Hua F, Bian Z. Coronavirus Disease 2019 (COVID-19): emerging and future challenges for dental and oral medicine. *J Dent Res* 2020;99:481–487
- ..Rational use of personal protective equipment for coronavirus disease 2019 (COVID-19). World Health Organization; February 27, 2020
- Zucco L, Levy N, Ketchandji D, et al. Perioperative Considerations for the 2019 Novel Coronavirus (COVID-19). Anesthesia Patient Safety Foundation. Available at: <https://www.apsf.org/news-updates/perioperative-considerations-for-the-2019-novel-coronavirus-covid-19/>. Accessed March 27, 2020
- Katzel EB, Basile P, Koltz PF, et al. Current surgical practices in cleft care: cleft palate repair techniques and postoperative care. *Plast Reconstr Surg* 2009;124:899–906
- Pradel W, Senf D, Mai R, et al. One-stage palate repair improves speech outcome and early maxillary growth in patients with cleft lip and palate. *J Physiol Pharmacol* 2009;60(suppl 8):37–41
- State of Florida. Emergency Management—COVID—19-Non-Essential Elective Medical Procedures. Available at: https://www.flgov.com/wp-content/uploads/orders/2020/EO_20-72.pdf. Accessed March 20, 2020
- Mississippi. State Department of Health COVID-19: Elective Surgical Procedures Must Be Rescheduled. https://msdh.ms.gov/msdhsite/_static/23,21854,341.html. Accessed March 19, 2020
- State of Alabama. Order of the State Health Officer Suspending Certain Public Gatherings Due to Risk of Infection by COVID-19. Available at: <https://governor.alabama.gov/assets/2020/04/Final-Statewide-Order-4.3.2020.pdf>. Accessed March 19, 2020
- State of Colorado. Ordering the Temporary Cessation of All Elective and Non-Essential Surgeries and Procedures and Preserving Personal Protective Equipment and Ventilators in Colorado Due to the Presence of COVID-19. Available at: https://www.colorado.gov/governor/sites/default/files/inline-files/D%202020%20009%20Ordering%20Cessation%20of%20All%20Elective%20Surgeries_0.pdf. Accessed March 19, 2020

18. State of Georgia. Gov. Kemp Issues New Executive Orders, Provides COVID-19 Update. Available at: <https://gov.georgia.gov/press-releases/2020-03-23/gov-kemp-issues-new-executive-orders-provides-covid-19-update>. Accessed March 23, 2020
19. State of Michigan. Temporary restrictions on non-essential medical and dental procedures. Available at: <https://www.michigan.gov/whitmer/0,9309,7-387-90499-522455-,00.html>. Accessed March 21, 2020
20. State of Minnesota. Directing Delay of Inpatient and Outpatient Elective Surgery and Procedural Cases during COVID-19 Peacetime Emergency. Available at: <https://www.leg.state.mn.us/archive/execorders/20-09.pdf>. Accessed March 19, 2020
21. State of Oregon. Conserving personal protective equipment and hospital beds, protecting health care workers, postponing non-urgent health care procedures, and restricting visitation in response to coronavirus (COVID-19) outbreaks. Available at: https://www.oregon.gov/gov/admin/Pages/eo_20-10.aspx. Accessed March 19, 2020
22. State of Vermont. Suspension of all Non-Essential Adult Elective Surgery and Medical and Surgical Procedures. Available at: <https://governor.vermont.gov/sites/scott/files/documents/ADDENDUM%203%20TO%20EXECUTIVE%20ORDER%2001-20.pdf>. Accessed March 20, 2020
23. State of Washington. Restrictions on Non Urgent Medical Procedures. Available at: <https://www.governor.wa.gov/node/495945>. Accessed March 19, 2020
24. DC Health. Recommendations on limitations of elective and non-urgent medical and dental procedures. Available at: https://dchealth.dc.gov/sites/default/files/dc/sites/doh/page_content/attachments/Letter%20-%20Elective%20Procedures.FINAL_.pdf. Accessed March 17, 2020
25. Riordan CP, Zurakowski D, Meier PM, et al. Minimally invasive endoscopic surgery for infantile craniosynostosis: a longitudinal cohort study. *J Pediatr* 2020;216:142–149.e2
26. Arko L, Swanson JW, Fierst TM, et al. Spring-mediated sagittal craniosynostosis treatment at the Children's Hospital of Philadelphia: technical notes and literature review. *Neurosurg Focus* 2015;38:E7
27. Noordzij N, Brouwer R, van der Horst C. Incomplete reossification after craniosynostosis surgery. *J Craniofac Surg* 2016;27:e105–108
28. Bruce WJ, Chang V, Joyce CJ, et al. Age at time of craniosynostosis repair predicts increased complication rate. *Cleft Palate Craniofac J* 2018;55:649–654
29. Louisiana Department of Health. Healthcare Facility Notice #2020-COVID19-ALL-06. Available at: <http://ldh.la.gov/index.cfm/newsroom/detail/5407>. Accessed March 18, 2020
30. North Carolina Department Of Health And Human Services. Available at: <https://www.ncdhhs.gov/divisions/public-health/covid19>. Accessed March 20, 2020
31. Ohio Department of Health. Re: Director's Order for the Management of Non-essential Surgeries and Procedures throughout Ohio. Available at: https://coronavirus.ohio.gov/wps/wcm/connect/gov/e7cee147-0f86-438b-ae1f-c5922f46c47c/Director%27s+Order+non-essential+surgery+3-17-2020.pdf?MOD=AJPERES&CONVERT_TO=url&CACHEID=ROOTWORKSPACE.Z18_M1HGGIK0N0JO00QO9DDDDM3000-e7cee147-0f86-438b-a. Accessed March 17, 2020
32. Society of Surgical Oncology. COVID-19 Resources. Available at: https://www.surgonc.org/resources/covid-19-resources/?_zs=eetRO1&_zl=DPqe5. Accessed March 24, 2020
33. Ueda M, Martins R, Hendrie PC, et al. Managing cancer care during the COVID-19 pandemic: agility and collaboration toward a common goal. *J Natl Compr Canc Netw* [published ahead of print March 20, 2020] doi: 10.6004/jnccn.2020.7560
34. Pennsylvania Department of Health. Guidance on ambulatory surgical facilities' responses to COVID-19. Available at: <https://www.health.pa.gov/topics/Documents/Diseases%20and%20Conditions/Guidance%20on%20Ambulatory%20Surgical%20Facilities%E2%80%99%20Responses%20to%20COVID-19.pdf>. Accessed March 20, 2020
35. State of Tennessee. An order to reduce the spread of COVID-19 by limiting non-emergency healthcare procedures. Available at: <https://publications.tnsosfiles.com/pub/execorders/exec-orders-lee25.pdf>. Accessed March 23, 2020

COVID-19 Pandemic and its Impact on Craniofacial Surgery

Brian T. Andrews, MD, MA,* Ravi Garg, MD,*
Wojciech Przylecki, MD,* and Mutaz Habal, MD†

Abstract: In late 2019, a novel coronavirus strain, SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2), also known as coronavirus disease 2019 (COVID-19), triggered a global pandemic as the virus spread from the Wuhan Province, China, across all continents. Although infrequent, severe respiratory infection and death caused by SARS-CoV-2 is disproportionately high amongst healthcare providers such as craniofacial surgeons who work in the head and neck region. Factors that impact SARS-CoV-2 transmission include: (1) high viral loads in the mucosa of the oral and nasopharynx, (2) limited and/or imprecise disease screening/confirmation testing, (3) access to and appropriate use of personal protective equipment (PPE).

Key Words: Coronavirus, SARS-CoV-2, COVID 19, severe acute respiratory syndrome coronavirus 2

In late 2019, the Wuhan Province of mainland China experienced an outbreak of a novel coronavirus strain, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), also known as coronavirus disease 2019 (COVID-19).¹ Similar to outbreaks of 2 other coronavirus pathogens, SARS-CoV and Middle East respiratory syndrome (MERS-CoV), SARS-CoV-2 is capable of causing severe and occasionally fatal (<10%) respiratory tract infection as it grows better in human airway epithelial cells than in standard tissue culture cells.^{2,3} In the subsequent months, the global spread of this viral pathogen has created a healthcare crisis not experienced since the Spanish Flu pandemic of 1918. As concerns for this disease progress, efforts to curb its progression in both the general population and in the healthcare professionals who treat these patients have been scrutinized. Healthcare providers who treat orofacial pathology such as craniofacial surgeons, otolaryngologists, dentists, oral surgeons, and ophthalmologists have an innately high risk of contracting SARS-CoV-2 compared with other providers.⁴ This is a direct result of an infected person's viral load being highest in the pharynx and upper aerodigestive tract.⁵

Zou et al further reported that asymptomatic carriers of SARS-CoV-2 can have positive viral cultures from the oro/nasopharynx for 7 to 11 days with no clinical signs of fever, cough, or shortness of breath. In a recent New England Journal of Medicine publication,

From the *Department of Plastic and Reconstructive Surgery, University of Kansas Medical Center, Kansas City, KS; and †Tampa Bay Craniofacial Center, Tampa Bay, FL.

Received March 31, 2020.

Accepted for publication March 31, 2020.

Address correspondence and reprint requests to Brian T. Andrews, MD, MA, Associate Professor, Department of Plastic and Reconstructive Surgery, University of Kansas Medical Center, Sutherland Institute, MS 3015, 3901 Rainbow Blvd, Kansas City, KS 66160; E-mail: bandrews@kumc.edu

The authors report no conflicts of interest.
Copyright © 2020 by Mutaz B. Habal, MD
ISSN: 1049-2275

DOI: 10.1097/SCS.00000000000006574